

The following citation list can be found on the **Partners in Information Access for the Public Health Workforce** Website at <http://phpartners.org/hp/>. This link is for the Healthy People 2010 Information Access Project section, where you will find a listing of many Healthy People 2010 focus areas, including vision. When you go to the vision page, you will be able to access National Library of Medicine PubMed searches related to each Healthy Vision 2010 objective. Below are citations for Objective 28-6. They were retrieved on February 29, 2004, limited to articles from the last 5 years.

Objective 28-6. Reduce visual impairment due to glaucoma.

Ahn BS, Kee C. Ability of a confocal scanning laser ophthalmoscope (TopSS) to detect early glaucomatous visual field defect. *Br J Ophthalmol.* 2000 Aug;84(8):852-5.

Allen CS, Sponsel WE, Trigo Y, Dirks MS, Flynn WJ. Comparison of the frequency doubling technology screening algorithm and the Humphrey 24-2 SITA-FAST in a large eye screening. *Clin Experiment Ophthalmol.* 2002 Feb;30(1):8-14.

Alward WL. Frequency doubling technology perimetry for the detection of glaucomatous visual field loss. *Am J Ophthalmol.* 2000 Mar;129(3):376-8.

Anton A. Early detection of glaucoma. *J Glaucoma.* 2003 Feb;12(1):59-62.

Arnold RW, Sitenga G. The detection of congenital glaucoma by photoscreen interpretation. *Alaska Med.* 2000 Jul-Sep;42(3):73-7.

Bach M. Electrophysiological approaches for early detection of glaucoma. *Eur J Ophthalmol.* 2001 Jul-Sep;11 Suppl 2:S41-9.

Bengtsson B. A new rapid threshold algorithm for short-wavelength automated perimetry. *Invest Ophthalmol Vis Sci.* 2003 Mar;44(3):1388-94.

Bobrow JC. Prospective intrapatient comparison of extracapsular cataract extraction and lens implantation with and without trabeculectomy. *Am J Ophthalmol.* 2000 Mar;129(3):291-6.

Booth AP, Anwar R, Chen H, Churchill AJ, Jay J, Polansky J, Nguyen T, Markham AF. Genetic screening in a large family with juvenile onset primary open-angle glaucoma. *Br J Ophthalmol.* 2000 Jul;84(7):722-6.

Bunce C, Hitchings R. Medical vs. surgical therapy in preventing visual field loss. *Ophthalmology.* 2003 Feb;110(2):249; author reply 249-50.

Burvenich H, De Clercq J. The combined IOP and CCT measurement in glaucoma screening. *Bull Soc Belge Ophthalmol.* 2000;276:15-8.

Coleman AL. Glaucoma. *Lancet.* 1999 Nov 20;354(9192):1803-10.

Corallo G, Gandolfo E. A method for detecting progression in glaucoma patients with deep, localized perimetric defects. *Eur J Ophthalmol.* 2003 Jan-Feb;13(1):49-56.

- Damji KF, Behki R, Wang L; Target IOP Workshop participants. Canadian perspectives in glaucoma management: setting target intraocular pressure range. *Can J Ophthalmol*. 2003 Apr;38(3):189-97.
- Distelhorst JS, Hughes GM. Open-angle glaucoma. *Am Fam Physician*. 2003 May 1;67(9):1937-44.
- Edwards JD, De Leon-Ortega J, Bearden WH, Rumble ME, Girkin CA. Is reading disability likely to interfere with glaucoma screening of adults using frequency-doubling technology perimetry? *Am J Ophthalmol*. 2003 Jun;135(6):816-20.
- Ellis JD, Morris AD, MacEwen CJ. Should diabetic patients be screened for glaucoma? DARTS/MEMO Collaboration. *Br J Ophthalmol*. 1999 Mar;83(3):369-72.
- Ellish NJ, Higginbotham EJ. Differences between screening sites in a glaucoma screening program. *Ophthalmic Epidemiol*. 2002 Oct;9(4):225-37.
- Ellish NJ, Higginbotham EJ. Evaluating a visual field screening test for glaucoma: how the choice of the gold standard affects the validity of the test. *Ophthalmic Epidemiol*. 2001 Dec;8(5):297-307.
- Erichev VP, Shamshinova AM, Egorova IV. [Psychophysical and electrophysiological tests in diagnosis of the early sign of glaucomatous optic neuropathy.] *Vestn Ross Akad Med Nauk*. 2003;(5):32-6. Russian.
- Fabre K, Michiels I, Zeyen T. The sensitivity and specificity of TOP, FDP, and GDX in screening for early glaucoma. *Bull Soc Belge Ophthalmol*. 2000;275:17-23.
- Faigen M. The early detection of glaucoma in general practice. *Aust Fam Physician*. 2000 Mar;29(3):282-5.
- Freeman PB. Recognizing limitations. *Optometry*. 2001 Apr;72(4):209.
- Fu P, Liu L, Li M, Yuan Y, Cui Y. [A clinical analysis of high myopia with primary open-angle glaucoma.] *Zhonghua Yan Ke Za Zhi*. 2002 Aug;38(8):480-3. Chinese.
- Garber N. Applanation tonometry. *J Ophthalmic Nurs Technol*. 1999 Nov-Dec;18(6):270-7; quiz 278-9.
- Gilchrist J. Optometric glaucoma referrals—measures of effectiveness and implications for screening strategy. *Ophthalmic Physiol Opt*. 2000 Nov;20(6):452-63.
- Goldberg I. Relationship between intraocular pressure and preservation of visual field in glaucoma. *Surv Ophthalmol*. 2003 Apr;48 Suppl 1:S3-7.
- Harper RA, Reeves BC. Glaucoma screening: the importance of combining test data. *Optom Vis Sci*. 1999 Aug;76(8):537-43.
- Harwerth RS, Smith EL 3rd, Chandler M. Progressive visual field defects from experimental glaucoma: measurements with white and colored stimuli. *Optom Vis Sci*. 1999 Aug;76(8):558-70.

Huang JT, Rhemtulla F, Huang PT. Glaucoma screening by primary care physicians in southern Alberta: patterns, methods, and deficiencies. *Can J Ophthalmol*. 2003 Jun;38(4):279-84.

Hyman L, Wu SY, Connell AM, Schachat A, Nemesure B, Hennis A, Leske MC. Prevalence and causes of visual impairment in The Barbados Eye Study. *Ophthalmology*. 2001 Oct;108(10):1751-6.

Iester M, Jonas JB, Mardin CY, Budde WM. Discriminant analysis models for early detection of glaucomatous optic disc changes. *Br J Ophthalmol*. 2000 May;84(5):464-8.

Johnson DH. Progress in glaucoma: early detection, new treatments, less blindness. *Ophthalmology*. 2003 Jun;110(6):1271-2.

Johnson DH. Progress in glaucoma: early detection, new treatments, less blindness. *Ophthalmology*. 2003 Apr;110(4):634-5.

Johnson LN. Medical vs. surgical therapy in preventing visual field loss. *Ophthalmology*. 2003 Feb;110(2):250-1.

Johnson LN. Glaucoma: our role in reducing the burden of blindness. *J Natl Med Assoc*. 2002 Oct;94(10):908-11.

Jonas JB, Bergua A, Schmitz-Valckenberg P, Papastathopoulos KI, Budde WM. Ranking of optic disc variables for detection of glaucomatous optic nerve damage. *Invest Ophthalmol Vis Sci*. 2000 Jun;41(7):1764-73.

Jonas JB, Budde WM. Diagnosis and pathogenesis of glaucomatous optic neuropathy: morphological aspects. *Prog Retin Eye Res*. 2000 Jan;19(1):1-40.

Kane W. Sight specific. *Nurs Times*. 1999 Aug 11-17;95(32):54-5.

King A, Migdal C. Clinical management of glaucoma. *J R Soc Med*. 2000 Apr;93(4):175-7.

Kipp MA. Childhood glaucoma. *Pediatr Clin North Am*. 2003 Feb;50(1):89-104.

Lee PP. Early detection of moderate glaucoma: redefining clinical care in 2001. *Arch Ophthalmol*. 2001 Jul;119(7):1069-70.

Lewandowska-Furmanik M, Matysik A, Pietras-Trzpiel M. The occurrence of primary open-angle glaucoma in a family. *Ann Univ Mariae Curie Skłodowska [Med]*. 2002;57(2):531-4.

Mackey DA, Craig JE. Predictive DNA testing for glaucoma: reality in 2003. *Ophthalmol Clin North Am*. 2003 Dec;16(4):639-45.

Martin SN, Sutherland J, Levin AV, Klose R, Priston M, Heon E. Molecular characterization of congenital glaucoma in a consanguineous Canadian community: a step towards preventing glaucoma-related blindness. *J Med Genet*. 2000 Jun;37(6):422-7.

Mills RP. Glaucoma screening as a vehicle for providing continuing medical education. *Acad Med*. 2000 Jan;75(1):91.

Mok KH, Lee VW. Nerve fiber analyzer and short-wavelength automated perimetry in glaucoma suspects: a pilot study. *Ophthalmology*. 2000 Nov;107(11):2101-4.

Nemesure B, Jiao X, He Q, Leske MC, Wu SY, Hennis A, Mendell N, Redman J, Garchon HJ, Agarwala R, Schaffer AA, Hejtmancik F; Barbados Family Study Group. A genome-wide scan for primary open-angle glaucoma (POAG): the Barbados Family Study of Open-Angle Glaucoma. *Hum Genet*. 2003 May;112(5-6):600-9. Epub 2003 Mar 04.

Nguyen RL, Raja SC, Traboulsi EI. Screening relatives of patients with familial chronic open-angle glaucoma. *Ophthalmology*. 2000 Jul;107(7):1294-7.

Palmisano P, Hynes M, Mueller L. Glaucoma and race: a case for screening in Connecticut. *Conn Med*. 2000 Feb;64(2):75-8. Erratum in: *Conn Med* 2000 Mar;64(3):188.

Parc CE, Johnson DH, Oliver JE, Hattenhauer MG, Hodge DO. The long-term outcome of glaucoma filtration surgery. *Am J Ophthalmol*. 2001 Jul;132(1):27-35.

Pieper B. National Glaucoma Awareness Month targets under-used Medicare screening benefit. *Optometry*. 2002 Dec;73(12):771-4.

Polo Llorens V, Pinilla Lozano I, Pablo Julvez LE, Larrosa Poves JM, Abecia Martinez E, Cuevas Andres R, Rojo Aragues A, Honrubia Lopez FM. [Short-wavelength automated perimetry (SWAP) in subjects with suspected glaucoma (I): assessment of sensitivity thresholds.] *Arch Soc Esp Oftalmol*. 2000 Feb;75(2):91-6. Spanish.

Quaranta GM, Vetrugno M, Trabucco T, Cantatore F, Cardia L. The role of frequency doubling technology (FDT) in glaucoma screening. *Acta Ophthalmol Scand Suppl*. 2000;(232):21-3.

Quigley HA, Jampel HD. How are glaucoma patients identified? *J Glaucoma*. 2003 Dec;12(6):451-5.

Remky A, Arend O. [Intra-individual variability of blue-yellow perimetry.] *Ophthalmologe*. 2000 Nov;97(11):774-80. German.

Rosenthal BP. Ophthalmology. Screening and treatment of age-related and pathologic vision changes. *Geriatrics*. 2001 Dec;56(12):27-31; quiz 32.

Schoevaerdts D, Levecq L, Vanpee D, Swine C. The medical management of primary open-angle glaucoma in the elderly patient. *Z Gerontol Geriatr*. 2003 Apr;36(2):121-3.

Simakova IL. [Visocontrastometry as one of diagnostic criteria for early detection of glaucoma and evaluation of stabilization of glaucomatous process.] *Vestn Oftalmol*. 2002 May-Jun;118(3):7-9. Russian.

Singer HW. Discam data offers early, objective way to study glaucoma. *J Ophthalmic Nurs Technol*. 2000 Mar-Apr;19(2):62-3.

Soderstrom M. Medical vs. surgical therapy in preventing visual field loss. *Ophthalmology*. 2003 Feb;110(2):249; author reply 249.

Stefan C, Nenciu A, Asandi R. [Primary open-angle glaucoma evolution after surgery.]

Oftalmologia. 2002;54(3):31-5. Romanian.

Tatemichi M, Nakano T, Tanaka K, Hayashi T, Nawa T, Iwasaki A, Miyamoto T, Hiro H, Sugita M. Laterality of the performance of glaucoma mass screening using frequency-doubling technology. *J Glaucoma.* 2003 Jun;12(3):221-5.

Tatemichi M, Nakano T, Tanaka K, Hayashi T, Nawa T, Miyamoto T, Hiro H, Iwasaki A, Sugita M; Glaucoma Screening Project (GSP) Study Group. Performance of glaucoma mass screening with only a visual field test using frequency-doubling technology perimetry. *Am J Ophthalmol.* 2002 Oct;134(4):529-37. Erratum in: *Am J Ophthalmol.* 2003 Sep;136(3):592.

Turkoski BB. Glaucoma and glaucoma medications. *Orthop Nurs.* 2000 Sep-Oct;19(5):71-6.

Vernon SA. Should diabetic patients be screened for glaucoma? *Br J Ophthalmol.* 1999 Sep;83(9):1096-7.

Viswanathan S, Frishman LJ, Robson JG, Walters JW. The photopic negative response of the flash electroretinogram in primary open-angle glaucoma. *Invest Ophthalmol Vis Sci.* 2001 Feb;42(2):514-22.

Vitale S, Smith TD, Quigley T, Kerrigan-Baumrind TA, Pease TE, Varma R, Friedman TS, Katz J, Tielsch JM. Screening performance of functional and structural measurements of neural damage in open-angle glaucoma: a case-control study from the Baltimore Eye Survey. *J Glaucoma.* 2000 Oct;9(5):346-56.

Wax MB, Camras CB, Fiscella RG, Girkin C, Singh K, Weinreb RN. Emerging perspectives in glaucoma: optimizing 24-hour control of intraocular pressure. *Am J Ophthalmol.* 2002 Jun;133 Suppl:S1-10.

Weinreb RN. A rationale for lowering intraocular pressure in glaucoma. *Surv Ophthalmol.* 2001 May;45 Suppl 4:S335-6.

Whitaker R Jr, Whitaker VB, Dill C. Glaucoma: what the ophthalmic nurse should know. *Insight.* 1999 Jul-Sep;24(3):86-91.

Wormald R. Treatment of raised intraocular pressure and prevention of glaucoma. *BMJ.* 2003 Apr 5;326(7392):723-4.

Xu L, Yang H, Zhao X. [A pilot study on rapid glaucoma screening.] *Zhonghua Yan Ke Za Zhi.* 2001 Jan;37(1):16-20. Chinese.

Yamada N, Chen PP, Mills RP, Leen MM, Stamper RL, Lieberman MF, Xu L, Stanford DC. Glaucoma screening using the scanning laser polarimeter. *J Glaucoma.* 2000 Jun;9(3):254-61.

Yamada N, Chen PP, Mills RP, Leen MM, Lieberman MF, Stamper RL, Stanford DC. Screening for glaucoma with frequency-doubling technology and Damato campimetry. *Arch Ophthalmol.* 1999 Nov;117(11):1479-84.

Yen MT, Gedde SJ, Flynn JT. Unilateral glaucoma in sotos syndrome (cerebral gigantism). *Am J Ophthalmol.* 2000 Dec;130(6):851-3.

[No authors listed] [Initiative for early diagnosis. Initiative to control glaucoma.] *Krankenpfl J*. 2003;41(4-6):119. German.

[No authors listed] Are you getting the screening tests you need? *Johns Hopkins Med Lett Health After 50*. 1999 Oct;11(8):4-5.